

## A record of *Thrichomys pachyurus* (Rodentia: Echimyidae) as prey of *Bothrops mattogrossensis* (Serpentes: Viperidae)

Ronildo Alves Benício<sup>1,\*</sup>, Karoline Ceron<sup>2</sup>, Matheus de Toledo Moroti<sup>3</sup>, and Diego José Santana<sup>4</sup>

Food is a crucial resource for all animals, and quantifying diets is one of the first steps in the study of species ecology (Sih and Christensen, 2001). What species eat and how and where this resource is obtained can help understand the use of resources and inter and intra-specific interactions (Granzinolli and Motta-Junior, 2007). However, the observation of predation records in nature is a fortuitous event (Huntingford, 1984). That is especially true for snakes, which are casually found during field works by active search (Fitch, 1987). Here, we present a record of predation event on a spiny rat *Thrichomys pachyurus* (Wagner, 1845) by the pitviper *Bothrops mattogrossensis* Amaral, 1925.

*Bothrops mattogrossensis* belongs to the *B. neuwiedi* complex, which is currently composed of eight species (Machado et al., 2014). *Bothrops mattogrossensis* occurs in Brazil, Peru, Bolivia, and Argentina (Silva and Rodrigues, 2008). In Brazil, the species occurs in the states of Amazonas, Rondônia, Mato Grosso do Sul, Tocantins, Goiás, and São Paulo (Silva and Rodrigues, 2008). Although considered as a terrestrial species (Martins et al., 2002), *B. mattogrossensis* is relatively slender and has the longest tail among other terrestrial

congeners, possibly an adaptation for climbing vegetation during seasonal floods of the open humid habitats in the Pantanal of western Brazil (Monteiro et al., 2006).

On 27 September 2017, in the dry season, at 13:45h, we found an adult female of *B. mattogrossensis* (Figure 1; Table 1) near the Base de Estudos do Pantanal (BEP), of the Universidade Federal de Mato Grosso do Sul (19.576297°S, 57.018474°W), municipality of Corumbá, state of Mato Grosso do Sul, Brazil. The individual was on the ground between the roots of a tree, on the border of a gallery forest, in a typical viper posture (Oliveira and Martins, 2001), and with part of the body expanded, indicating presence of undigested (or partially digested) stomach content (Figure 1A). We collected the individual and, after necropsy, the stomach content was removed, which was identified as being a subadult male (body mass = 120 g; see Andreazzi et al., 2011) of *Thrichomys pachyurus* (Wagner, 1845) (Figure 1B). The fact that the anterior and posterior limbs were pointed backwards indicates that the rat was eaten by the head first (see Cabral et al., 2017). *Thrichomys pachyurus* is a large rodent species belonging to the Echimyidae family (Braggio and Bonvicino, 2004). This species has a wide distribution (Lacher, 2016) and occurs in Paraguay, Bolivia and central and western Brazil in the states of Tocantins, Goiás, and Mato Grosso (Pessoa et al., 2015). The rodent represents 22% of the total length of the snake, and 61% of its weight (Table 1). We used the same measurements presented in Cabral et al. (2017). Both specimens were housed in Coleção Zoológica da Universidade Federal de Mato Grosso do Sul (*B. mattogrossensis* = ZUFMS-REP 02628, and *T. pachyurus* = ZUFMS-MNV 00059).

*Bothrops mattogrossensis* does not show an ontogenetic shift in diet from ectothermic to endothermic prey and, compared to the other species of the *B. neuwiedi* complex (e.g., Martins et al., 2002; Valdujo et al., 2002;


<sup>1</sup> Programa de Pós-Graduação em Ecologia e Recursos Naturais, Universidade Federal de São Carlos, São Carlos, São Paulo, Brazil.

<sup>2</sup> Programa de Pós-Graduação em Ecologia e Conservação, Universidade Federal do Mato Grosso do Sul, Campo Grande, Mato Grosso do Sul, Brazil.

<sup>3</sup> Programa de Pós-Graduação em Biologia Animal, Universidade Federal do Mato Grosso do Sul, Campo Grande, Mato Grosso do Sul, Brazil.

<sup>4</sup> Departamento de Zoologia, Universidade Federal do Mato Grosso do Sul, Campo Grande, Mato Grosso do Sul, Brazil.

\* Corresponding author. E-mail: benicio.ufscar@gmail.com



**Figure 1.** An adult female of *Bothrops mattogrossensis* with part of the body expanded, indicating presence of undigested (or partially digested) stomach content (A), from municipality of Corumbá, state of Mato Grosso do Sul, Brazil. The same specimen dissected with its prey, a subadult male of *Thrichomys pachyurus* (B). The position pointed backwards of anterior and posterior limbs indicates that the rat was eaten by the head first (see Cabral *et al.*, 2017). A calliper with 15 cm long was used as a scale to the photos.

Hartmann *et al.*, 2004), has a generalized diet, preying small mammals, anurans, lizards, snakes, and centipedes (Monteiro *et al.*, 2006). Both traits are considered plesiomorphic for the genus *Bothrops* (Martins *et al.*, 2002). This is the first record of predation of *Thrichomys pachyurus* – a large species of rodent (individuals can reach 700 g of body mass; Pessôa *et al.*, 2015).

Predation events with relatively large-sized prey have been recorded for some species of the genus *Bothrops* (e.g., Cabral *et al.*, 2017; Cadena-Ortiz *et al.*, 2017). Some authors have suggested that the relation between

prey size and snake size follows the efficient foraging theory, which assumes that bigger preys would be preferred because they maximize net energy or nutrient gain and minimize the time and energy spent acquiring a quantity of resources (Sih and Moore, 1990). Furthermore, it has been demonstrated that females of the *B. neuwiedi* complex have larger body sizes and proportionally larger heads than males, which may reflect sexual differences in diet (Martins *et al.*, 2002; Valdujo *et al.*, 2002; Hartmann *et al.*, 2004). Larger head and body size of females are a possible adaptation

**Table 1.** Measurements in centimetres of the individuals of *Thrichomys pachyurus* (ZUFMS-MNV 00059) and *Bothrops mattogrossensis* (ZUFMS-REP 02628) from municipality of Corumbá, state of Mato Grosso do Sul, Brazil.

Body measurements	<i>T. pachyurus</i>	<i>B. mattogrossensis</i>
Snout-vent length	14.3	87.3
Tail length	6.7	5.2
Total length	21	92.5
Head width	2.8	24.4
Head height	2.9	16.1
Body width	7.1	25.9
Body height	2.65	23.4
Weight (in grams)	238	390

for ingesting larger prey items (Pough and Groves, 1983; Shetty and Shine, 2002). However, there is no evidence supporting sexual differences in the diet of *B. mattogrossensis* (Monteiro et al., 2006).

For Downes (2002), patterns of size-selective predation by snakes are determined by differential prey vulnerability rather than by active predator choice. Regarding *B. mattogrossensis*, further studies on prey availability and vulnerability associated to dietary analyses may help understand the prey composition of this species.

**Acknowledgements.** This article is result of a research conducted during the 20th Curso de Campo do Pantanal (EcoPan2017) of the Universidade Federal de Mato Grosso do Sul. We thank Grazielle C.G. Soaresini for help in snake necropsy; Luiz Gustavo Oliveira-Santos for identifying the rodent species. Ronildo A. Benicio thanks Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, Proc. 2015/11821-0) for financial support. Karoline Ceron and Matheus de T. Moroti thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior for scholarships.

## References

- Andreazzi, C.S.D., Rademaker, V., Gentile, R., Herrera, H.M., Jansen, A.M., D'andrea, P.S. (2011): Population ecology of small rodents and marsupials in a semi-deciduous tropical forest of the southeast Pantanal, Brazil. *Zoologia* **28**: 762–770.
- Braggio, E., Bonvicino, C.R. (2004): Molecular divergence in the genus *Thrichomys* (Rodentia, Echimyidae). *Journal of Mammalogy* **85**: 316–320.
- Cabral, H., Rojas, V., Galluppi, T., Ortiz, E., Baez, M. (2017): Comments on the diet of *Bothrops alternatus* Duméril, Bibron and Duméril, 1854. *Herpetology Notes* **10**: 219–220.
- Cadena-Ortiz, H., Barahona-V., A., Bahamonde-Vinueza, D., Brito, M.J. (2017): Anecdotal predation events of some snakes in Ecuador. *Herpetozoa* **30**: 93–96.
- Downes, S.J. (2002): Size-dependent predation by snakes: selective foraging or differential prey vulnerability? *Behavioral Ecology* **13**: 551–560.
- Granzinolli, M.A.M., Motta-Junior, J.C. (2007): Feeding ecology of the White-tailed Hawk (*Buteo albicaudatus*) in south-eastern Brazil. *Emu* **107**: 214–222.
- Hartmann, M.T., Marques, O.A.V., Almeida-Santos, S.M. (2004): Reproductive biology of the southern Brazilian pitviper *Bothrops newwiedi pubescens* (Serpentes, Viperidae). *Amphibia-Reptilia* **25**: 77–85.
- Huntingford, F.A. (1984): Some ethical issues raised by studies of predation and aggression. *Animal Behaviour* **32**: 210–215.
- Fitch, H.S. (1987): Collecting and life-history techniques. In: *Snakes: Ecology and Evolutionary Biology*, p. 143–164. Seigel, R.A., Collins, J.T., Novak, S.S. Eds., New York, USA, MacMillan Publishing Company.
- Lacher, T. (2016): *Thrichomys pachyurus*. The IUCN Red List of Threatened Species 2016: e.T136245A22206322. Available at: <http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T136245A22206322.en>. Accessed on 07 June 2018.
- Machado, T., Silva, V.X., Silva, M.J. (2014): Phylogenetic relationships within *Bothrops newwiedi* group (Serpentes, Squamata): geographically highly-structured lineages, evidence of introgressive hybridization and Neogene/Quaternary diversification. *Molecular Phylogenetic Evolution* **71**: 1–14.
- Martins, M., Marques, O.A.V., Sazima, I. (2002): Ecological and phylogenetic correlates of feeding habits in Neotropical pitvipers of the genus *Bothrops*. In: *Biology of the Vipers*, p. 307–328. Schuett, G.W., Höggren, M., Douglas, M.E., Greene, W.H., Eds., Eagle Mountain, USA, Eagle Mountain Publishing.
- Monteiro, C., Montgomery, C.E., Spina, F., Sawaya, R.J., Martins, M. (2006): Feeding, reproduction and morphology of *Bothrops mattogrossensis* (Serpentes, Viperidae, Crotalinae) in the Brazilian Pantanal. *Journal of Herpetology* **40**: 408–413.
- Oliveira, M.E., Martins, M. (2001): When and where to find a pitviper: activity patterns and habitat use of the lancehead, *Bothrops atrox*, in central Amazonia, Brazil. *Herpetological Natural History* **8**: 101–110.
- Pessôa, L.M., Tavares, W.C., Neves, A.C.A., Silva, A.L.G. (2015): Genus *Thrichomys*. In: *Mammals of South America, Volume 2, Rodents*, p. 989–999. Patton, J.L., Pardiñas, U.F.J., D'elía, G.D., Eds., Chicago, USA, University of Chicago Press.
- Pough, F.H., Groves, J.D. (1983): Specialization of the body form and food habits of snakes. *American Zoologist* **23**: 443–454.
- Shetty, S., Shine, R. (2002): Sexual divergence in diets and morphology in Fijian sea snakes *Laticauda colubrina* (Laticudinae). *Austral Ecology* **27**: 77–84.
- Sih, A., Moore, R. (1990): Interacting effects of predator and prey behavior in determining diets. In: *Behavioural mechanisms of food selection*, p. 771–796. Hughes, R., Ed., Heidelberg, DE, Springer-Verlag.
- Sih, A., Christensen, B. (2001): Optimal diet theory: when does it work, and when and why does it fail? *Animal Behaviour* **61**: 379–390.
- Silva, V.X., Rodrigues, M.T. (2008): Taxonomic revision of the *Bothrops newwiedi* complex (Serpentes, Viperidae) with description of a new species. *Phyllomedusa: Journal of Herpetology* **7**: 45–90.
- Valdujo, P.H., Nogueira, C., Martins, M. (2002): Ecology of *Bothrops newwiedi pauloensis* (Serpentes: Viperidae: Crotalinae) in the Brazilian Cerrado. *Journal of Herpetology* **36**: 169–176.